

**IBUPROFEN- ibuprofen tablet, coated**  
**Discount Drug Mart**

-----  
**Drug Mart 44-291**

***Active ingredient (in each brown tablet)***

Ibuprofen USP,  
200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
- menstrual cramps
- toothache
- the common cold
- backache
- headache
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- skin reddening
- asthma (wheezing)
- rash
- facial swelling
- shock
- blisters
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - leg swelling
  - slurred speech
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

***Other information***

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

***Inactive ingredients***

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

***Questions or comments?***

**Call 1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

***Principal display panel***

DISCOUNT  
**drug**  
**mart**

FOOD FAIR

NDC 53943-291-15

MAY CONTAIN  
**ANTI-THEFT**  
DEVICE

**IBUPROFEN**  
**TABLETS USP, 200 mg**

*Pain Reliever / Fever Reducer (NSAID)*

Actual Size

Compare to the active ingredient in ADVIL<sup>®</sup> Tablets†

**50 COATED TABLETS**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS  
BROKEN OR MISSING**

†This product is not manufactured or distributed by Wyeth LLC, owner of the registered trademark Advil<sup>®</sup> Tablets.

50844 REV0618A29115

**DISTRIBUTED BY: DRUG MART - FOOD FAIR**  
**211 COMMERCE DRIVE, MEDINA, OH 44256**  
**[www.discount-drugmart.com](http://www.discount-drugmart.com)**

---

**SATISFACTION GUARANTEED**


IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE  
TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO  
STORE, SEND REASON FOR DISSATISFACTION, NAME,  
ADDRESS AND EMPTY PACKAGE TO:

DISCOUNT DRUG MART  
211 COMMERCE DRIVE, MEDINA, OHIO 44256

---

IBUPROFEN			
ibuprofen tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-291
Route of Administration	ORAL		
Active Ingredient/Active Moiety			

No print/No varnish  
Lot & Exp date




0 9335102769 7

**IBUPROFEN**  
TABLETS USP, 200 mg  
Pain Reliever / Fever Reducer (NSAID)

Compare to the active ingredient in ADVIL® Tablets!

Actual  
Size



50 COATED TABLETS

**Drug Facts** (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - leg swelling
  - slurred speech
- weakness in one part or side of body

**Drug Facts** (continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

**Inactive ingredients:** canauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate,

**Questions or comments?**  
Call 1-800-426-9391 8:30 AM-4:00 PM ET.  
Monday-Friday

**Drug Facts** (continued)

microcrystalline cellulose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

**Drug Facts**

**Keep Outer Package for Complete Product Information**

**Active Ingredient**  
(in each brown tablet)  
Ibuprofen USP  
200 mg (NSAID) \*nonsteroidal anti-inflammatory drug

**Purpose**

**Uses**

- temporarily relieves minor aches and pains due to:
  - menstrual cramps
  - toothache
  - the common cold
  - headache
  - muscular aches
  - backache
  - minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- rash
- skin redness
- facial swelling
- asthma (wheezing)
- shock
- blisters
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you take more or for a longer time than directed.

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

**Drug Facts** (continued)

microcrystalline cellulose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?**  
Call 1-800-426-9391 8:30 AM-4:00 PM ET.  
Monday-Friday

**Drug Facts** (continued)

This product is not manufactured or distributed by Wyeth LLC, owner of the registered trademark Advil®

50844 REV0618A29115  
DISTRIBUTED BY: DRUG MART - FOOD FAR  
211 COMMERCE DRIVE, MEDINA, OH 44256  
www.discount-drugmart.com

**Drug Mart 44-291**

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	

Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;291
<b>Contains</b>			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-291-15	1 in 1 CARTON	05/24/1988	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:53943-291-12	1 in 1 CARTON	05/24/1988	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:53943-291-14	500 in 1 BOTTLE; Type 0: Not a Combination Product	05/24/1988	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/24/1988	

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(53943-291)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	pack(53943-291)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	pack(53943-291)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(53943-291)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(53943-291)

Revised: 4/2021

Discount Drug Mart